

TABLE OF CONTENTS

1. Introduction
2. Water Scarcity and Avocado Production in California
3. Analytical Framework
4. Analytical Model
5. Empirical Approach
6. Results
7. Conclusion and Policy Implications
8. References
9. Appendix

**Dealing with Water Scarcity:
Adoption of Water Saving Technologies and Management Practices
by California Avocado Growers**

Julie Reints¹, Ariel Dinar² and David Crowley¹

¹Department of Environmental Sciences, ²School of Public Policy, University of California, Riverside, CA 92521

Summary:

The irrigated agriculture sector has been facing an increased scarcity in terms of both quantity and quality of water worldwide. Consequently, the sustainability of water intensive crops, such as avocado, is threatened when water becomes too expensive, or growers must shift to saline water supplies that reduce crop yields. A variety of irrigation technologies and water saving management practices are now recommended to help growers through times of limited water supplies. To examine how growers adopt different practices and combinations of practices, we used Kohonen self-organizing maps and developed logit models to identify the most common bundles of technologies and management practices that growers are using to deal with water scarcity. We test the validity of the proposed bundles and factors affecting their adoption using primary data obtained from a survey of California avocado growers at the heights of the drought during 2012-2013. Results show that in selecting bundles, or in adopting single technologies or management practices, farm location, share of income from agricultural production, use of cooperative extension advice, and farmer characteristics such as age and education, all play important roles in grower adoption of methods to adapt to water scarcity.

Acknowledgements: The authors would like to acknowledge financial support from the California Avocado Commission via grant No. 12034311; from the Giannini Foundation of Agricultural Economics, Mini Grant Fund. Julie Reints would like to acknowledge the support from the Graduate Student Dissertation Mentorship Fellowship, from the Herbert Kraft Scholarship, Campbell Research Fellowship Award, and from the Dean's Distinguished Fellowship. Ariel Dinar would like to acknowledge support from the NIEA Multistate Project W3190 "Management and Policy Challenges in a Water-Scarce World."

Dealing with Water Scarcity:

Adoption of Water Saving Technologies and Management Practices by California Avocado Growers

Julie Reints¹, Ariel Dinar² and David Crowley¹

¹Department of Environmental Sciences, ²School of Public Policy, University of California,
Riverside, CA 92521

Keywords: adoption—avocado—bundles—California—irrigation technology—water conservation—water scarcity

JEL codes: Q33, Q16, Q25

Acknowledgements: The authors would like to acknowledge financial support from the California Avocado Commission via grant No. 12034311; from the Giannini Foundation of Agricultural Economics, Mini Grant Fund. Julie Reints would like to acknowledge the support from the Graduate Student Dissertation Mentorship Fellowship, from the Herbert Kraft Scholarship, Campbell Research Fellowship Award, and from the Dean’s Distinguished Fellowship. Ariel Dinar would like to acknowledge support from the NIFA Multistate Project W3190 “Management and Policy Challenges in a Water-Scarce World.”

Authors Bio: Julie Reints is a PhD Graduate, Department of Environmental Sciences. David Crowley is Emeritus Professor, Department of Environmental Sciences. Ariel Dinar is a Distinguished Professor, School of Public Policy, University of California, Riverside, USA.

Dealing with water scarcity:

Adoption of water saving technologies and management practices by California avocado growers

Abstract: The irrigated agriculture sector has been facing an increased scarcity in terms of both quantity and quality of water worldwide. Consequently, the sustainability of water intensive crops, such as avocado, is threatened when water becomes too expensive, or growers must shift to saline water supplies that reduce crop yields. A variety of irrigation technologies and water saving management practices are now recommended to help growers through times of limited water supplies. To examine how growers adopt different practices and combinations of practices, we used Kohonen self-organizing maps and developed logit models to identify the most common bundles of technologies and management practices that growers are using to deal with water scarcity. We test the validity of the proposed bundles and factors affecting their adoption using primary data obtained from a survey of California avocado growers at the heights of the drought during 2012-2013. Results show that in selecting bundles, or in adopting single technologies or management practices, farm location, share of income from agricultural production, use of cooperative extension advice, and farmer characteristics such as age and education, all play important roles in grower adoption of methods to adapt to water scarcity. [192 words]

Introduction

Worldwide availability of good quality water for irrigated agriculture is significantly affected by climate change and increased urban and agricultural demands. Extended periods of drought further exacerbate the already dwindling stocks and flows of existing ground and surface water. During periods of drought, growers are faced with a need to find solutions to sustain production and meet the world's demand for sustenance.

Growers may respond to lower water availability and reduction in quality (e.g., higher salinity) by introducing various water conservation technologies and management practices with short- and long-term implications. They may alter their irrigation scheduling by taking advantage of monitoring and scheduling programs that are available in the market; they may follow part of

their land and adjust the irrigation area to the available water that they have been allocated. In the case of orchard crops such as avocado, farmers may switch to alternate more drought tolerant crops such as lemon, prune the trees to reduce the canopy size, or “stump” the trees to remove the canopy and hold production for one or more years until they can again receive regular quantities of irrigation water. During drought, farmers also intensify their consultation with experts, including agricultural extension agents and advisors representing agricultural input firms. Lastly, farmers may invest in changes to their irrigation technologies, soil water monitoring equipment, new wells, and pipelines for use of alternative water sources such as treated wastewater.

All these, and other responses can be undertaken by growers either separately or jointly, as bundles of responses. Bundling, or combining technologies, takes place when growers use several technologies and management practices that support each other instead of adopting one technology or management practice independently. Adoption of bundles may provide growers more flexibility than adoption of individual technologies or management practices¹ (Fleischer et al. 2011).

The existing literature has dealt with factors that affect decision making and adoption of agricultural technologies for many years, in many countries and under many different physical and institutional settings. While irrigation technology adoption in general and in California,² in particular, has been studied in the literature (e.g., Campbell and Dinar, 1993; Dinar and Yaron, 1992; Feder et al. 1985; Feder and Umali, 1993; Koundouri et al. 2006), few works have been published on “bundling” water-saving technologies and management practices that conserve water in agriculture (e.g., Fleischer et al. 2011; Wang et al. 2010). Wang et al. (2010) examined the strategy of using different cropping patterns (crop bundles) with different water requirements by

¹ From hereafter we use the terms irrigation technologies, management practices/strategies, and conservation techniques interchangeably.

² Our paper focuses on adoption of water saving technologies and management practices by avocado growers in Southern California, as will be explained and justified in the next section.

Chinese farmers and found that, depending on the region, certain crop bundles provide farmers flexibility in dealing with climate change impacts on water scarcity. Fleischer et al. (2011) found that Israeli growers bundle their crop mix, and irrigation and crop cover technologies in response to changes in long-term availability of water. When given the opportunity to choose from a variety of options they choose to bundle their agricultural technologies (cover, irrigation technologies, crop mix) in order to adapt to increased water scarcity, providing flexibility and sustainability across topography and climate.

Such flexibility in response may be needed for agricultural growers facing limiting climatic conditions such as those faced by California growers located along the state coast, from north to south, where significant differences in rainfall and temperature occur. Bundling water technologies in order to adapt to change in climate and water scarcity provide resiliency and results in higher profits, as was observed in Wang et al. (2010) and Fleischer et al. (2011).

So far, there are several gaps in research on agricultural technology adoption. Most studies (e.g., Feder et al. 1985, Dinar and Yaron 1992, Koundouri et al. 2006, Tey and Brindal 2012) identify a single type of technology and determine its adoption likelihood and pattern by growers. Two examples are irrigation methods and precision agricultural technologies. But irrigation technologies themselves may not be sufficient for increasing water efficiency. Irrigation technology such as drip irrigation and micro sprinklers are typically treated as a separate decision from fertilization and precision agricultural technologies in adoption and diffusion rates (Feder et al., 1985; Feder and Umali, 1993; Koundouri et al., 2006). Scientific evidence (Sureshkumar et al., 2017) suggests that bundling drip or micro-sprinkler irrigation with fertilization (Fertigation) leads to much more effective results.

Other aspects not considered in existing adoption studies include the use of water saving technologies and management practices that can help growers guide irrigation practices. These are practices and technologies aimed at reducing water use while at the same time maintaining yields and increased profit from water savings. Soil moisture monitoring technologies have been shown to significantly increase the water use efficiency of avocado trees (Kiggundu et al., 2012). Growers use irrigation calculators, deficit irrigation, and water shortage mitigation practices to save on-farm water use. These strategies, also known as sustainable irrigation management practices or irrigation best management practices, include technologies and practices that are resource conserving and are presented as a variety of options available to growers (Boland et al., 2006; Pereira et al., 2002).

We focus in this work on Avocado growers in California. Based on Carmen (2019) Avocado, mainly Hass variety (99.4%), is grown in California on nearly 50,000 bearing acres (as of 2017) by nearly 5,000 growers. Avocado gained popularity in the USA due to its healthy attributes. Avocado is a water intensive crop, sensitive to water quantity and its salinity. With a water consumption of 4 acre-feet (AF)/acre and price charged by water utilities, ranging between \$1200-1300 per AF during periods of water scarcity (UCCE and CAS, 2012), avocado growers in Southern California have to consider additional responses. In particular, the 2010-2016 prolonged drought led some of them introduce drastic responses to the drought, including use of desalinated groundwater by mobile treatment units to increase supply, and/or stumping of entire plots to temporarily reduce water demand.

Currently, there is no published work on avocado production and the determinants of adoption with respect to water saving practices or response to decreasing water supplies and qualities. What makes some growers adopt advanced water saving technologies and management

practices while other growers do not is an important policy question. The present study covers aspects of adoption that have not been addressed in previous research on adoption of water saving technologies and management practices. The paper specifically aims to identify water technologies and water conservation techniques that are available to avocado growers in California for conservation of water; how those choices are bundled and how selected socioeconomic and farm characteristics contribute to adoption.

We start with our methodology, followed by the analytical model used along with the empirical approach. Then we describe the data collection process, and the bundle determination methodology. We then present results from the empirical models that were estimated. We wrap up with conclusions and the policy implications of the findings. The paper includes auxiliary appendixes. We review water saving technologies and management practices used by California avocado growers in Appendix 1. And Appendix 2 presents the Kohonen Self-Organizing Maps.

Methodology

Based on existing research we developed an analytical framework with expected effects of the key variables that can explain adoption of water conserving technologies and practices. Once the basis for our framework was established, we moved in two directions: First, we constructed an analytical model, based on the framework, which will be used for the statistical estimates for examining our hypotheses on criteria that affect adoption of selected technologies alone or when combined in bundles. Second, we developed and administered a questionnaire to collect primary data in the regions we planned to focus on. Finally, we specified functional forms to be estimated in order to verify the hypotheses.

Analytical Framework

We include in our analytical framework several factors that have been used in the literature to explain the extent of adoption of agricultural technologies in the context of irrigation water use. Based on the literature, economic factors such as cost of water, farmer characteristics (education, experience and age);³ farm characteristics factors such as farm location, soil properties and landscape, farm size, share of farm income from agriculture, and farm management structure; and informational factors such as sources of know-how, all contribute to adoption of irrigation technologies. A detailed description is provided below for selected attributes, and a summary of these variables and their associated expected effects on water technology and management practices adoption is provided in Table 1.

Table 1 about here

Socio-economic Factors: Farmer Characteristics

Previous studies on adoption of technologies have shown that human capital is a critical factor in identifying adopters. Koundouri et al. (2006) found that under production uncertainty human capital played an important role in adoption of technologies in Crete where younger, more educated farmers were adopters of water efficient irrigation technologies. In studies estimating single technology and bundled technology adoption, education has been a common determinant for predicting adoption, where the higher the level of education the greater is the extent of adoption (Dorfman 1996; Wang et al., 2010; Fleischer, et al., 2011; Robertson et al., 2012).

Use of bundles doesn't necessarily mean more complexity. In many cases bundles actually imply simplicity and improved precision of the agronomic process. The case of combining irrigation and fertilization (fertigation) is an example for simplification and improvement of the

³ As opposed to the a-priori belief age and experience are not necessarily correlated. For example, the USDA-ERS (2012) study found that 35 percent of beginning (unexperienced) farmers are over age 55 and nearly 13 percent are 65 or older. We found that among the avocado growers in our study in California, the correlation between years of age and years of experience is 0.323 and not significant.

precision in the combined process compared to the irrigation and fertilization when done separately. Education could be more important in certain bundles. For example, Fleicher et al. (2011, Table 2), and Wang et al. (2010, Table 1) find that education is not significant in explaining selection of certain bundles.

When compared to less educated farmers in precision fertilizer technologies, higher farmer education level also resulted in faster adoption by growers (Feder and Umali, 1993). Genius et al. (2014) found that the combination of age and education had an effect on adoption where younger educated growers adopted drip technology. Grower's age and formal education is considered in our paper as well (we do not measure speed of adoption in this paper).

Experience is an important factor in adapting to climate change. In a study that considered choice of crop and bundled technologies, growers who had more farming experience choose to grow a crop mix that includes an orchard crop, which is considered more profitable than row crops (Fleischer, et al., 2011). Hence, we can propose that years of experience may influence adoption in our context as well.

Agro-ecological Factors: Farm Characteristics

Regions with high aridity along with a sandy soil can increase water inputs and consequently increase crop production risks, especially in the event of drought (Koundouri et al., 2006). Many studies have shown that local weather is an important factor in adoption, where farms with higher evapotranspiration (ET_o) rates and high aridity indices adopt water saving technologies (Dinar and Yaron 1992; Campbell and Dinar 1993; Koundouri et al., 2006).

Several studies have determined that farm size is important when considering technology adoption. Farm size can greatly influence adoption as larger farms may have access to higher

equity and monetary resources to invest in water saving equipment (Dorfman 1996; Koundouri et al., 2006; Wang et al. 2010).

Complexity in operating the irrigation system may be an important impediment to adoption. Soil has the ability to retain moisture based on its unique texture, organic matter and these physical properties are not uniformly distributed throughout an orchard (Saxton and Rawls 2006, Farmer et al., 2003). For example, a soil type of a plot located on hill slopes may have different physical properties than a soil type of a plot located on leveled ground. This sub-field variability in soil properties, shape of the irrigation plot, and topography add to an orchard's irrigation complexity, which represents the level of difficulty a farmer experiences in setting up and maintaining an irrigation system. We propose that the higher the irrigation complexity a grower faces, the harder it is for the grower to manage water resources and make decision about water management and thus will reduce the likelihood of adopting new technologies and practices..

Informational Factors

Recently, Genius et al. (2014) investigated the role of information transmission in promoting agricultural adoption and diffusion through extension services and social learning with olive growers in Greece. They found that farmers are primarily informed about technologies by their interactions with agricultural extension services and other farmers. Tiarniyul (2009) developed the notion of a technology score to measure technology advancement, and found that a farmer's technology score was affected significantly by number of extension visits, in addition to years of formal education, farming experience and land ownership status, when they examined adoption of technologies among rice farmers in Nigeria. Informational factors can be considered both where growers seek information to manage their orchards and what topics are sought (Genius et al., 2014; Tey and Brindal, 2012). Not only where growers seek information is important, but the extent of

use of that avenue of information (Genius et al., 2014); more adoption will occur among growers who see information collected from these sources as being useful (Tey and Brindal, 2012). In addition to extension, growers obtain information and knowhow also from commercial providers of inputs, professional associations, neighbors, and others.

Analytical Model

Avocado growers can respond to water shortages and climate change in several ways. Assuming profit-maximizing behavior, and given that water is one of the highest cost components in avocado production in California, growers will choose as many technologies and management practices that will decrease water consumption while keeping a high profit. Therefore, we estimate a model that explains the selection of a technology (or a bundle of technologies) such that the profit of the grower is maximized, subject to the conditions the grower faces.

Based on the work reviewed in the previous section, we introduce two models to capture the behavioral relationship of choosing a bundle of irrigation technologies and water management practices in avocado production. First, we estimate a model where we look at adoption of either irrigation technology and/or management practices in a binary-choice framework as affected by a set of variables such as farm characteristics, farmer characteristics, informational variables and fixed effects variables.⁴ Second, we estimate a model of selection of bundles of technology and management practices also explained by the same set of independent variables as in the first model. The fixed effects in our models include county location, which captures attributes that are associated with the county and have not been captured by the other variables. We use a cross section data to explain what affects at a given time period—the midst of the California drought,

⁴ Fixed effect picks up any variation in the dependent variable that happen over time (for time/year fixed effects for example) or over space/location (for country or city fixed effects for example) and that is not attributed to the other explanatory variables. Fixed effects are estimated usually by introducing dummy variables that represent each year or each location (in the examples we used).

avocado growers choice of a single technology or management practice, or bundles of technologies and management strategies in order to save water.

The profit function (we show only the multinomial bundle model because the logit and multinomial model are similar in notation), adapted from Fleischer et al. (2011), for choosing a bundle is:⁵

$$Y_j = Y_j(r, m, k) + \varepsilon_j, \quad j = 1, 2, \dots, J$$

where J is the total number of technologies and management bundles to control water use. Y_j is a dichotomous function (0-1) indicating whether or not bundle j is selected. Y_j is a function of a vector of farmer characteristics, r , a vector of farm characteristic, m , (including climatic effects embedded in location), and a vector of informational factors, k . A farmer will choose a bundle j iff $Y_j > Y_i \quad \forall j \neq i$, thus we can say that the probability of a farmer choosing bundle j is:

$$P_j = \Pr(Y_j > Y_i) \quad \forall j \neq i$$

A critical assumption of the multinomial logit procedure is that the relative probability of any two alternative bundles is not affected by adding a third bundle.

Assuming that ε is independently Gumbel distributed and the profit function can be written linearly in its parameters, as $Y_j = \alpha + \beta r_j + \gamma m_j + \delta k_j$, where α, β, γ , and δ are the estimated coefficients, then the probability, P_j , can be calculated as follows:

$$P_j = \frac{e^{\alpha + \beta r_j + \gamma m_j + \delta k_j}}{\sum_{i=1}^J e^{\alpha + \beta r_i + \gamma m_i + \delta k_i}}$$

where P_j is the probability that a given bundle, comprising of technologies and management practices, will be selected.

⁵ A logit model estimates the probability of dichotomous range for the dependent variable, such as yes/no, pass/fail, measured as 1/0 as a function of a set of explanatory variables. A multinomial logit model estimates the probabilities of different possible outcomes of a categorically distributed dependent variable, given a set of independent variables.

The analytical framework will be tested using two model sets. The first model set is a logit model where a grower's adoption of any technology or management practice with regards to water management in their orchard is analyzed as a binary decision. We developed two logit versions, using two different sets of explanatory variables in each. We followed Dinar et al. (1992) who referred to "modern irrigation technologies" that include any of the following: solid-set sprinklers, micro-sprinklers and drip. Some technologies or management practices may need higher skills (such as more experience, higher education, more support from extension, and higher financial resources for purchasing and for operation of the technology) than others. By referring to "any tech/practice" with assumed equal sophistication we may have ended up with an over-estimation of some of the coefficients, such as of education and experience, which could limit the generalization of our results.

The second model set is a multinomial logit model that combines eight water saving technologies and managements into bundles of likely technologies used in combination by the growers. Each grower in the analysis was assigned to a bundle based on their use of the technologies and management practices. Four bundles (0, 1, 2, 3) were used in a logit regression with non-users (bundle 0) being the benchmark to which coefficients are compared.

We turn now to the data and the constructed variables we use in the statistical estimates.

Data and Variable Construction

Farmer and farm information is primary data that we obtained directly from avocado growers. Data was collected using a survey instrument comprised of 71 questions that was administered during 2012-2013 (See Appendix 3 in the Auxiliary Material). The survey was distributed to California

avocado growers with the help of the California Avocado Commission (CAC),⁶ using its database of growers. Surveys were distributed by email, mail and in face-to-face interactions during growers' meetings. Responses were received from 128 growers. Five responses were not used due to data quality issues. Of the remaining 123 responses, 94, 17, and 12 were obtained by email, mail, and direct interviews in grower meetings, respectively. Since the information received was not subjective but rather fact-based, we are not concerned of possible sample bias.

The 123 observations represent a total of 3899 acres⁷ of avocado orchard, nearly 7 percent of the avocado farmland in California. The total response accounts for nearly 2 percent of the number of avocado growers in California (distribution by county can be found in Table 2). Location of growers that responded in each California County is presented in Figure 1.

One could argue that the dataset exhibits selection bias because it could be claimed that large growers volunteered their time in answering the survey questions. However, selection bias has been ruled out because the distribution of avocado farm size in the sample follows a similar distribution pattern as in actual avocado farms within each county in which avocado is grown in California (Table 2).

Figure 1 about here

Table 2 about here

Water Saving Technologies Used in the Empirical Models

Growers in our sample use different technologies to maintain or increase profits and address risk management due to water availability fluctuations. We identified (Escalera et. al., 2015) methods

⁶ CAC organizes all Avocado growers in California for advocacy, promotion and knowledge creation. All producing acres are levied a tax per unit of yield to allow the operation of CAC.

⁷ 1 acre = 0.4 hectares.

of water saving technologies and management practices and have used them as a starting point for categorizing the bundles.

The Water saving technologies and management practices reported by the growers are listed below:

Soil moisture measuring devices (Allowing growers to monitor and maintain adequate soil moisture in real time, determine when irrigation is needed, and when sufficient water has been applied): (1) Soil auger, (2) tensiometer, (3) gypsum block, (4) dielectric sensors, (5) capacitance/dielectric sensors, (6) neutron probe (7) gravimetric.

Irrigation calculators (Designed to help growers determine site specific crop water requirements based on weather data and crop coefficients): (1) CIMIS—California Irrigation Management Information System that uses transpiration rates to determine crop specific water needs on a daily basis.

Water saving techniques: (1) pruning (2) stumping, (3) removing trees, (4) turning off water to trees to reduce on farm water use, (5) any tree management to reduce water use.

Water management techniques: (1) improving Distribution Uniformity with the use of water audits, (2) testing irrigation water for salinity management, (3) test soil moisture with sensors, (4) measure soil salinity (5) irrigating by calendar (instead of using monitoring equipment or CIMIS).

Irrigation Technologies: (1) pressure compensating sprinklers, (2) micro-sprinklers, (3) Drip irrigation.

Miscellaneous techniques: (1) Choose district water over groundwater, when possible, (2) use “By Feel” method to decide when to irrigate. The “By Feel” method is a simple way to assess the level of moisture in the soil using hand feel.

Setting the Bundles

A total of 23 water saving technologies and management practices were identified from the administered survey. In order to explore the relationships between the variables, the data was first analyzed using Kohonen Self-Organizing Maps (KSOM) (Kohonen 2013) generated with the software program Synapse (Peltarion, Inc., Stockholm). KSOM uses an artificial neural network to reduce the dimensionality of highly dimensional data sets into two dimensional arrays that allow visualization of the relationships between all of the variables of interest. In the map, each variable is assigned a grid or tile that is color coded to illustrate the range of values that are observed in the data set (Kohonen 2001). Similarities in distributed color patterns represent correlations across all of the dependent and independent variables, such that variables with similar patterns can be designated as potentially belonging to a cluster. KSOM methods are especially useful for comparing non-parametric data, and have been widely used in many scientific applications. Examples of prior applications related to agriculture include land use classification, examination of relationships between farm practices and water quality, and factors associated with farm profitability (see review in Kalteh et al., 2008). Here we applied this approach to identify individual variables that behaved similarly and that could thus be grouped into hypothetical bundles for further statistical analysis and modeling of factors affecting adoption of these technologies for dealing with water scarcity.

The concept of bundling is based on the idea that individual growers could use more than one technology and water management practice to respond to changes in water availability. Likewise, some technologies were little used or were sporadically adopted. For example, the SOM shows no apparent relationship between variables describing grower characteristics and the use of the gravimetric methods, neutron probe, or dielectric and capacitance sensors for monitoring soil water availability. As a result, those technologies were eliminated from the analysis. We were also

able to exclude drip and micro sprinkler irrigation technologies since almost all growers have these irrigation technologies. Also, since only some growers have both the choice of groundwater or district (surface) water whereas other growers cannot select among these sources, we excluded whether the use of groundwater or district water affected adoption. We should also indicate that technological and management practices bundles are both location (country, region) specific and technology and managing practice specific (in Wang et al, 2011 they are a combination of crops, and in Fleischer, et al., 2011 they are a combination of crops and technologies).

Based on the KSOM analysis (Appendix 2), technologies and practices were narrowed down to fewer choices. Similar patterns shown in the SOM tiles represent associations between particular dependent and independent variables in the dataset. Out of original 23 water savings and management practices reported and answered by growers, 8 were selected as the basis for the bundles used in the estimation of the adoption models: (1) water audit, (2) Soil moisture by feel, (3) Soil moisture by gypsum block, (4) Soil moisture by tensiometer; (5) Irrigation using calendar, (6) Irrigation using CIMIS, (7) Management of tree canopy, and (8) pressure compensating sprinklers. In this study, growers may bundle up to 8 discrete types of technologies or management practices to conserve water.

Zero Bundle Growers

Bundle zero, the default, or the benchmark outcome, represents no adoption of practices or technologies, or a set of practices and technologies other than those reported in the questionnaire and identified by SOM, that are used by a group of growers to determine when and how much to irrigate. For instance, growers adopting bundle zero may have irrigation system infrastructure limits where avocado is irrigated along with other crops since they cannot separate the two. The orchard may be irrigated by a management company and owners do not know when irrigation

events take place or do not have control over how decisions are made with respect to water management. Also, part time growers who are able to only irrigate when they are physically present fall in this category as they can turn on the water only when they can visit the orchard (Faber, 2015).

Development of other Bundles

To determine the most likely combination of technologies and managements practices used by the growers, a multiple correspondence analysis (MCA) was used to identify the most commonly grouped selections by growers (Figure 2). Since growers can choose up to 8 different methods, in any combination available, the MCA proved to be a useful tool to identify likely combinations that growers use in practice.

MCA is a multivariate analysis that conceptually is similar to Principal Component Analysis (PCA), but applies to categorical rather than continuous data (Linting et al., 2002). PCA is a statistical procedure that uses an orthogonal transformation in order to convert a set of observations that could have correlated variables (and thus create a problem in residing in the right hand side as explanatory variables in a regression) into a set of linearly uncorrelated variables that could be presented as one variable (Jolliffe, 2012). Both PCA and MCA provide a means of displaying or summarizing a set of data in, simpler, two-dimensional graphical form. This type of analysis can be used to detect underlying structures in the dataset.

Correlations between variables shown in the MCA coordinate graph (Figure 1) accounted for 72.5% of the inertia (extent) in the adoption of technologies/management practices. The higher the total inertia in the dimensions of the technologies, the better is the model fit. The first and second dimensions combined accounted for 79.5% variance in the dataset. The results of the MCA presented in Figure 1 show the first two dimensions plotted against each other and distributed in

four quadrants that delineate a correlation between variables. Each variable is shown in four quadrants with the associated binary response, 1= Yes, 0= No.

The multiple correspondence graph shows that in the upper left quadrant, growers who answered ‘yes’ to: irrigating by calendar, using pressure compensating sprinklers, tensiometers and used “by feel” method are correlated with each other. In the upper right hand quadrant, growers who answered ‘no’ to utilizing water audits, ‘no’ to using CIMIS, and ‘no’ to stumping or heavily pruning to conserve water are correlated with each other. In the lower left hand quadrant, growers who answered ‘yes’ to having to stump or heavily prune to conserve water, ‘yes’ to getting water audit to improve water efficiency, ‘yes’ to using CIMIS and ‘yes’ to using gypsum blocks are correlated with each other. In the lower right-hand quadrant, growers who answered ‘no’ to using tensiometers, ‘no’ to using “by feel”, ‘no’ to irrigating by calendar, and ‘no’ to using pressure compensating sprinklers are correlated with each other.

Figure 2 about here

The Bundles in the Dataset

Bundle 0: This bundle includes growers who do not use any of the 8 water management methods described earlier.

Bundle 1: This bundle represents growers who use pressure compensating sprinklers, by feel method, tensiometers and irrigation by calendar. This bundle is the least advanced, as it requires the least amount of training, education and funding to use. Although tensiometers require knowledge of water soil relations, they are inexpensive and easy to use.

Bundle 2: This bundle represents growers who have had to stump or heavily prune their trees to conserve water, use CIMIS, gypsum blocks, and utilize water audits to improve on-farm water use efficiency. This bundle is more sophisticated compared to bundle 1. Using CIMIS, although free,

requires knowledge of evapotranspiration concepts and learning how to use the model with respect to seasons and type of crop. Utilizing water audits requires knowledge of irrigation systems, how to improve the water efficiency and willingness to pay for improvements after the audit is completed.

Bundle 3: This bundle represents growers who use a combination of technologies and management methods from Bundle 1 and Bundle 2 to include: pressure compensating sprinklers, by feel, tensiometers, calendar-based irrigation, stump/prune trees, CIMIS, gypsum blocks, and water audits. This bundle is the most flexible in use and may represent growers that need flexibility in how they approach water management. Bundle 3 is the most sophisticated bundle.

Table 3 presents the distribution of the bundles within the dataset of study growers. Sixteen percent of growers are associated with bundle 0. Bundle 2 and bundle 3, constitute twelve and thirty five percent of the sample, respectively. Bundle 3 constitute thirty seven percent of the sample, including growers that use a combination all eight of the technologies and management practices identified in the study. These growers did not fit into a single category as found in the MCA and were assigned a bundle category of 3 so they could be represented in this study.

Table 3 about here

Based on the literature review, we can hypothesize that more sophisticated bundles such as 2 and 3 will be adopted by growers that have higher education, value cooperative extension and have a high share income from avocado production.

Explanatory Variables

We used information collected in our survey to create the set of explanatory variables. We distinguish between *Farm Acres* and *Avocado Acres* where some growers may specialize only in avocado and some may have mixed cropping systems in which avocado is interplanted with other

trees such as lemon or cherimoya. Of the 6 avocado growing counties in the sample we found that the coastal locations are similar and different as a group than the inland counties. Therefore, we used Ventura, San Luis Obispo, Santa Barbara, and Orange counties as benchmark and created 2 dummy variables: *Riverside County* and *San Diego County*. Additional variables in our analysis are *Irrigation Complexity* (index), *Agricultural Water Rate* (\$/hcf), *Age of owner* (years), *Formal Education* (ranking), *Years' Experience* (years), *Share Income from Avocado*, *Original Owner* (0/1), *Use of Cooperative Extension* (ranking), *Leaf Sampling* (0/1), *Follow Lab Recommendation* (0/1), *Test Irrigation Water* (0/1). All of the explanatory variables fall into categories that represent farm characteristics, farmer characteristics and informational factors as described in Table 1.

Empirical models

We developed 2 sets of empirical models. The first set included 2 logit equations (f_1 and f_2) that explain the likelihood of selecting any technology or management practice by the farmers. The left hand-side of the equation is a dichotomous variable (0/1) having a value 1 if there is any use of a water saving technology and or management practice, and a value of 0 otherwise. On the right hand-side of the estimated equations we included two different sets of explanatory variables that were explained and justified in our Analytical Framework Section. Model f_1 includes farm-level, informational, and human capital variables. Model f_2 includes in addition the regional fixed effects variables. The third model (g_i) that we estimated is the multinomial logit regression model where we explained in a system of equations estimated simultaneously the likelihood of selecting the 3 bundles with reference to bundle 0 (as a benchmark). The multinomial logit model includes all farm-level, human capital, informational, and fixed regional effects variables.

The specific empirical models we estimated and the dependent and independent variables included in each of the models are provided below:

Logit (dichotomous) regression: selecting a technology/management practice = f_1 (grow crops other than avocado, avocado acres, cost of irrigation water, owner age, education, share income from avocado, cooperative extension use)

Logit (dichotomous) regression: selecting a technology/management practice = f_2 (owner operated, farm acres, riverside county, San Diego county, irrigation complexity, owner age, education, share income from avocado, cooperative extension use)

Multinomial logit regression: selecting bundle $i = g_i$ (owner operated, farm acres, Riverside county, San Diego county, irrigation complexity, owner age, education, share income from avocado, cooperative extension use).

Several variables are suspected of showing endogeneity: Bundle adoption decision could be expected to be jointly determined with share income from avocado, with level of education, and with use of extension. It could well happen that unobservable factors other than these regressors could affect the decision to select any of the bundles. To check for a possible endogeneity in our model we employed the Durbin–Wu–Hausman (DWH) test, or the augmented regression test for endogeneity (Hausman, 1978). The Hausman test for endogeneity result (6.90; 0.648) failed to be rejected, indicating that there is no endogeneity and no need for instrumentation.

Results

We start with descriptive results and then turn to the results of the econometric estimates.

Descriptive Statistics Results

The average grower manages over 3000 trees with a weighted average (depending on variety) of 137 trees per acre. Growers grow on different soil textures (sandy, loam, clay), and face various irrigation block shapes (rectangular, square, irregular shaped), and topography (% grade) on each

irrigation block they manage. Growers manage anywhere from 1 to 20 (or more) irrigation blocks depending on the size of their orchard, age of trees, topography, or water delivery. These variables were aggregated, using the MCA to represent the irrigation complexity a grower is faced with in managing water. Typically, irregular shaped irrigation blocks are harder to manage with respect to irrigation systems and blocks on steep slopes face non-uniform water delivery challenges. These variables were assigned categorical numbers from least complex to most complex with respect to water management and then aggregated to calculate a variable that accounts for irrigation complexity. The higher the value of the aggregated variable the more complex is the irrigation of the orchard.

Data on water source and quality was collected from the growers. Water quality data from the growers was checked against public records provided by water districts. Data on groundwater quality was provided by Groundwater Monitoring and Assessment Program (GAMA), Geotracker of USGS. Average chloride concentration for all surface water sources, districts, and groundwater sources, is 68.7 mg L⁻¹, TDS is 552 mg L⁻¹ and Electric Conductivity (EC) is 0.76 dS/m.⁸ Avocado is mainly irrigated with district water (82%), though some growers have access to both surface and groundwater sources and can irrigate the orchard by mixing water supplies in a pond or by using different water sources at different times during the growing season. Irrigation technology includes micro-sprinklers (87% of the orchards) or drip (7% of the orchards), with very few growers that use another type such as overhead sprinklers (up to 3% of the orchards). The average age of the micro-sprinklers and drip irrigation was 15 years and 2.3 years respectively.

⁸ Long term irrigation with water having an EC > 1 dS m⁻¹ is detrimental to avocado production. The ability of avocado roots to take up water from the soil via osmosis is reduced and leads to water stress and closure of the leaf stomata. Use of saline water also commonly leads to reductions in yield caused by chloride toxicity when chloride is present in the irrigation water at concentrations greater than 100 mg L⁻¹. Avocado is considered to be one of the most sensitive of all crops to soil salinity and requires very careful irrigation management to avoid soil salinization when using high EC irrigation water supplies (UCCE and CAS, 2012).

An average avocado grower in our survey manages 93 acres of farmland, of which 31 acres are in avocado production. This suggests that the remaining acres are used for other crops, fallowed, or with buildings. Indeed, when asked if growers grow crops other than avocado 65% responded that they grow another crop such as citrus, grapes, persimmons, olives and ornamental crops. The average age of a grower in our sample was 62 years and they were mostly male (80%). Farm management is mostly in sole proprietorship (64.4%) and next, in partnerships (21%). A high percent of growers had formal higher education, either graduate degrees (43.8%) or Bachelor's degrees (33.9%). Growers had an average of 19.7 years of experience growing avocado, and 42 percent of the growers in the sample were original owners of the orchards they currently manage. Growers reported earning as much as 20 percent of their overall income from avocado production.

Our sample suggests the following distribution (%) of growers among California counties: 41, 28, 12, 10, 6, and 3 in San Diego, Ventura, Riverside, San Luis Obispo, Santa Barbara, and Orange counties, respectively. Our sample suggests also land distribution among these counties to be 26, 41, 19, 4, 8, and 2, respectively (Table 2). The counties of San Diego and Riverside capture 69% and 67% of the growers and avocado land, respectively. Both the distribution of our sample growers and land across the counties reflect their actual distribution (Table 2).

While we use only Cooperative Extension as a source of information and knowhow, avocado growers indicated that their sources include also the California Avocado Commission, journals, suppliers and other growers.

Descriptive statistics of the variables that are included in our econometric analyses is presented in Table 4. Riverside and San Diego counties were used in the regressions as dummies and Orange, San Luis Obispo, Santa Barbara and Ventura were aggregated and used as a

benchmark for several reasons. First, we found that Riverside and San Diego counties were the most arid avocado growing regions and it could be inferred that these counties are more affected by droughts and lack of high-quality irrigation water. These preliminary observations are in line with previous work (e.g., Dinar et al., 1992; Genius et al, 2014; Fleischer et al., 2011), suggesting that in our case facing harsher climatic conditions, or higher levels of scarcity, would lead avocado growers in these regions to be more likely adopters of water saving technologies and management practices. Second, when the models were run we found collinearity between Orange, San Luis Obispo, Santa Barbara and Ventura Counties suggesting that there are similarities between these counties that should be considered in the empirical model.

Table 4 about here

We also tested for possible multi-collinearity among the variables that we use to explain the likelihood of selecting a technology or management practice in the 2 logit models f_1 and f_2 and the likelihood to select bundle i in the multinomial logit model g_i . We found no correlation between pairs of variables that were used in the right hand-side of the same estimated equation. Some of the variable pairs that were correlated were not included in the same regression. Therefore, we exclude the possibility of multicollinearity in our models.

Regression Results

Data were analyzed with both logit regression (Table 5) and multinomial logit regression (Table 6) models. The logit model predicts the likelihood of adopting a specific technology and the multinomial logit predicts the likelihood of adopting a given bundle. Two discrete logit models (Table 5) were used to estimate adoption of any irrigation technology and/or any management practice as a dichotomous variable (0/1). A multinomial model was used when considering

adoption as a bundled choice, with non-bundle-adopters (bundle 0) used as benchmark and bundles 1-3 as explained earlier (Table 6).

Two logit regressions were chosen that consider adoption as a binary choice, adopter versus non-adopters, taking into account farm characteristics, farmer characteristics and informational factors (Table 5). The difference between these two models are that exogenous factors such as farm characteristics that are modified to test the theory that farm location in arid counties has an effect on adoption while holding all other variables constant. The multinomial logit regression uses explanatory variables that are also used in the logit regression to compare between adopters and non-adopters.

Based on Model 1 (Table 5) the probability of being an adopter of irrigation managements and/or technologies increases when orchards are located in Riverside County, when growers have a higher income share from avocado production, and when growers are provided information by cooperative extension. Riverside County has the highest aridity index in the sample and is located in the inland desert region of the CIMIS evapotranspiration map. Regression results of Model 1 suggests that the probability of being an adopter decreases with irrigation complexity and owner's age. The results of Model 1 are consistent with previous work and with our expectations.

We estimated the impact of additional exogenous farm factors in a second logit regression model (Model 2, Table 5) in which the farm characteristics included growing other crops, and also considered avocado acres and cost of water. In this model, we find that probability of choosing to adopt increases with share income from avocado and with cooperative extension support, and decreases with owner's age. These results are consistent with previous work and with our expectations.

For the multinomial logit regression, the most significant factors affecting the selection of bundle 1 (Table 6) were share income from avocado production (mean income was 16%) and informational factors such as use of cooperative extension. We did not find that farm characteristic, such as location, had an important contribution to selecting bundle 1.

We found that farm characteristics and location of orchard, were important for a growers to choose bundle 2 with Riverside County being statistically significant. Farmer characteristics such as age, education and share of income from avocado explained decision to select bundle 2. In addition, use of cooperative extension was significant in selecting bundle 2.

The probability of a grower using Bundle 3, where growers could choose any combination and weren't limited to a discrete selection, was increased by farm location in both San Diego and Riverside County, the most arid regions of California. Riverside and San Diego Counties are the most southern, warmer and drier counties where avocado is grown in the state. Owner's age decreased the probability of using bundles 2 and 3. This trend has been seen in previous related literature (Genius et al., 2014). The probability of selecting bundle 3 was decreased by the farm's irrigation complexity. Income share from avocado and use of cooperative extension were also important factors affecting a grower's decision to select bundle 3 for irrigation management.

While the logit models in Table 5 cannot be fully contrasted with the multinomial logit model in Table 6, still we can draw some comparisons. Both logit and multinomial logit estimated have their significant coefficients with similar signs, which suggest a robustness across these models. Among the fixed regional effects, Riverside county is positive and significant in the logit (model 1 where it appears) and in bundles 2 and 3 of the multinomial logit model. San Diego county is positive and significant only in bundle 3 of the multinomial logit model. As we know, the aridity index of Riverside county is much higher than that of San Diego county, which can

explain its insignificance up to the level of sophistication of bundle 3. These coefficients measure the influence of water scarcity on the decision to adopt the water technologies and management practices. The variable irrigation complexity, measuring the difficulty facing growers in adopting new technologies, is negative and significant in the logit (model 1 where it appears) and in bundle 3 of the multinomial logit model, most likely due to the higher specifications imposed by bundle 3. The negative coefficient is interpreted as the inhibitive effect of complicated landscape on adoption. The coefficient of the age variable is negative and significant in all logit models and in bundles 2 and 3 of the multinomial logit model. The coefficient of age in the equation of bundle 1 is insignificant, suggesting that the adoption of a relatively simple bundle is not affected by age. Education is not significant in most estimates except in bundle 2 of the multinomial logit model. We do not have a good explanation for that finding and the fact it is not significant in our models (while in many previous work it was found to be positive and significant) except for speculating that we may have encountered measurement errors in the way education was recorded. Share income from avocado was found positive and significant in all models and this is expected in our hypotheses and observed in previous studies. Finally, cooperative extension was found also highly significant and positive, which, again is expected and in agreement to previous work. In summary, our results of the two sets of models provide consistent results that are also consistent with previous finding in the literature.

Conclusion and Policy Implications

The research concluded in this paper identifies a set of variables that affect the adoption of water conservation technologies and management practices by avocado growers in California. One of the contributions of this research is the identification and adoption of bundles of technologies and

management practices rather than adoption of a single technology or management practice. The results of the analysis in this study support previous attempts to look at adoption as a multi-faceted activity motivated by resource scarcity, economics factors, physical determinants, and information provision.

One of the most important findings implies that informational factors such as cooperative extension have an important role in adoption of water saving technologies and management practices for California avocado growers. Cooperative extension agents are able to distribute research and tools necessary for growers to mitigate the impact of drought. Growers gain knowledge by meeting cooperative extension specialists in avocado tree management during visits to the farm, regional meetings, office hours, or phone calls, and obtain information via publications made available to the general public. Cooperative extension was found to be positive and significant in all estimated equations, concluding that it is an important source of information for growers.

The research also shows that human capital variables, in particular age, are important in predicting how a grower makes decisions about water management. Economics of avocado production, measured as share of income from avocado in the farm was found to be an important factor influencing grower adoption of any individual technology or bundles. This implies that when facing water scarcity and if avocado is an important source of farm income a grower is more likely to adopt water management practices to sustain profitable production.

One important conclusion from our research, realized through the multinomial logit model is that growers will need to have more flexibility in their approach to water management to mitigate climate change and reductions in irrigation water quantity and quality. Growers who were able to select from many different discrete management tools to manage water were located in Riverside

and San Diego counties and had less complex irrigation systems. Riverside and San Diego Counties have higher aridity indexes and are predominantly on district water, typically a more expensive option for growers. In areas where there is less water available to growers, they may benefit from simplifying their irrigation systems in order to facilitate maintenance and improve water-use efficiency. The need for increased flexibility in technology adoption under increased water scarcity goes hand in hand with the role of cooperative extension. Therefore, future importance of cooperative extension services to the avocado production industry is amplified as future climate change impacts are expected to worsen.

Policy Implications

There are several policy implications that emerge from this paper. First, we found that a combination of policy interventions should be considered to support farmers that face water scarcity, starting with creating and making information available by extension agents, designing policies that address farms with different farm irrigation complexities, water sources, human capital levels (education, age, experience), and farm management structures.

While the differences between the two counties San Diego and Riverside were said to be the result of the aridity, it can be also that differences in county services play an important role in enabling farmers adoption of single technologies and bundles. A policy that acknowledges these additional services could make a difference in farmers' success when facing water scarcity and deteriorated quality.

We realize that there are differences among regions and thus, policy responses should take into account regional differences and provide a quilt rather than an umbrella policy. Regions with large urban centers could also take advantage of an additional source of water in the form of treated wastewater that may benefit farmers if properly treated and adequately priced.

There are a couple of caveats that need to be addressed. First, our sample size may affect the ability to capture differences in responses of farmers to scarcity, beyond what we were able to capture. While our sample is representative in terms of number and land distribution across the various counties, we may have not captured the actual number of technologies used. A second caveat that has to be mentioned is the possible bias from assuming that all technologies require a similar set of human capital skills when adopted separately (in the logit analyses). While we follow the practice used in the literature, attributing similar complexities and skill needs to a range of technologies and management practices (defined as “any technology or management practice”), still this may not be sufficient and the adoption of each technology or management practice should be estimated specifically. With a larger number of observations this could be performed in a more statistically significant manner.

References

- Ascough, G. W. and G. A. Kiker (2002). The effect of irrigation uniformity on irrigation water requirements. *Water SA*, 28, 235-241.
- Boland, A. M., D. Bewsell and G. Kaine (2006). Adoption of sustainable irrigation management practices by stone and pome fruit growers in the Goulburn/Murray Valleys, Australia. *Irrigation Science*, 24, 137-145.
- Campbell, M. B. and A. Dinar (1993). Farm organization and resource use. *Agribusiness (New York)*, 9, 465-480.
- California Avocado Commission (2013). <http://www.californiaavocado.com/>
- Carmen, H. F. (2019). The Story Behind Avocados' Rise to Prominence in the United States. *ARE Update*, 22(5):9-11. University of California, Giannini Foundation of Agricultural Economics.
- Dinar, A. and D. Yaron (1992). ADOPTION AND ABANDONMENT OF IRRIGATION TECHNOLOGIES. *Agricultural Economics*, 6, 315-332.
- Dinar, A., M. B. Campbell, and D. Zilberman (1992). Adoption of Improved Irrigation and Drainage Reduction Technologies Under Limiting Environmental Conditions. *Environmental & Resource Economics*, 2:373-398.
- Dorfman, J. H. (1996). Modeling multiple adoption decisions in a joint framework. *American Journal of Agricultural Economics*, 78, 547-557.
- Escalera, J., A. Dinar and D. Crowley (2015). Adoption of Water-Related Technology and Management Practices by the California Avocado Industry. *Agricultural and Resource Economics Update: Giannini Foundation of Agricultural Economics*, 18(3):5-8.
- Faber, B. (2015). Personal Communications.
- Farmer, D., M. Sivapalan and C. Jothityangkoon (2003). Climate, soil, and vegetation controls upon the variability of water balance in temperate and semiarid landscapes: Downward approach to water balance analysis. *Water Resources Research*, 39, 21.
- Feder, G., R. E. Just and D. Zilberman (1985). ADOPTION OF AGRICULTURAL INNOVATIONS IN DEVELOPING-COUNTRIES - A SURVEY. *Economic Development and Cultural Change*, 33, 255-298.
- Feder, G. and D. L. Umali (1993). The Adoption of Agricultural Innovations - A Review. *Technological Forecasting and Social Change*, 43, 215-239.

- Ferreya, R., Maldonado, P., Celedón, J., Gil, P. M., Torres, A., and G.Selles (2008). Soil Air Content Effects on The Water Status of Avocado Trees. *ISHS Acta Horticulturae* 792, 291-296.
- Fleischer, A., R. Mendelsohn and A. Dinar (2011). Bundling agricultural technologies to adapt to climate change. *Technological Forecasting and Social Change*, 78, 982-990.
- Genius, M., P. Koundouri, C. Nauges and V. Tzouvelekas (2014). Information Transmission in Irrigation Technology Adoption and Diffusion: Social Learning, Extension Services, and Spatial Effects. *American Journal of Agricultural Economics*, 96, 328-344.
- Gil, P. M., Bonomelli, C., Schaffer, B., Ferreyra, R., & Gentina, C. (2012). Effect of soil water-to-air ratio on biomass and mineral nutrition of avocado trees. *Journal of Soil Science and Plant Nutrition*, 12(3), 609-630.
- Green, S. R., A. Hodson, M. Barley, M. Benson and A. Curtis (2012). Crop IR Log - an Irrigation Calculator for Tree and Vine Crops. *Viii International Symposium on Sap Flow*, 951, 277-284.
- Hofshi, R. e. a. (2010). Yearbook of the California Avocado Society for the year 2010. 93, 51-71.
- Hausman, J. A. (1978). Specification Tests in Econometrics. *Econometrica*. 46 (6), 1251–1271.
- Jolliffe, I.T. (2002). *Principal Component Analysis*, second edition, Berlin: Springer.
- Kalteh, A.M., P. Hjorth, R. Berndtsson (2008). Review of the Self-organizing Map (SOM) Approach in Water Resources: Analysis, Modelling and Application. *Environmental Modelling & Software*, 23:835-845.
- Kiggundu, N., K. W. Migliaccio, B. Schaffer and J. H. Crane (2012). Water Savings, Nutrient Leaching, and Fruit Yield in a Young Avocado Orchard as Affected by Irrigation and Nutrient Management. *Irrigation Science*, 30(4):275-286.
- Kohonen, T. (2011). *Self-organizing Maps*. Berlin: Springer (2001).
- Kohonen, T. (2013) Essentials of the self-organizing map. *Neural Networks*, 37, 52-65.
- Koundouri, P., C. Nauges and V. Tzouvelekas (2006). Technology adoption under production uncertainty: Theory and application to irrigation technology. *American Journal of Agricultural Economics*, 88, 657-670.
- Kozlowski, T. T. (1997). Responses of Woody Plants to Flooding and Salinity. *Tree Physiology*, 17(7):490, doi:10.1093/treephys/17.7.490 Key: citeulike:13604913

- Linting, M., Meulan, J. J., Groenen, P. J. F. and van der Kooij, A.J. (2007). Nonlinear Principal Components Analysis: Introduction and Application. *Psychological Methods* 12(3): 336-358.
- Oster, J. D. and M. L. Arpaia (2002). Setting TMDLs for salinity and chloride based on their effects on avocado (Hass) productivity. *USCID Water Management Conference on helping irrigation agriculture adjust to TMDLs (Total maximum daily loads - for water quality), Sacramento, California, USA, 23-26 October 2002*, 241-252 ref.
- Pereira, L. S., T. Oweis and A. Zairi (2002). Irrigation management under water scarcity. *Agricultural Water Management*, 57, 175-206.
- Robertson, M. J., R. S. Llewellyn, R. Mandel, R. Lawes, R. G. V. Bramley, L. Swift, N. Metz and C. O'Callaghan (2012). Adoption of variable rate fertiliser application in the Australian grains industry: status, issues and prospects. *Precision Agriculture*, 13, 181-199.
- Saxton, K. E. and W. J. Rawls (2006) Soil water characteristic estimates by texture and organic matter for hydrologic solutions. *Soil Science Society of America Journal*, 70, 1569-1578.
- Schaffer, B., B. N. Wolstenholme & A. W. Whiley (2013). *The Avocado: Botany, Production and Uses*.
- Stagakis, S., Gonzalez-Dugo, V., Cid, P., Guillen-Climent, M. L., and Zarco-Tejada, P. J. (2012). Monitoring Water Stress and Fruit Quality in an Orange Orchard under Regulated Deficit Irrigation Using Narrow-band Structural and Physiological Remote Sensing Indices. *ISPRS Journal of Photogrammetry and Remote Sensing*, 71, 47-61. doi:10.1016/j.isprsjprs.2012.05.003.
- Sureshkumar, P., Geetha, M.C. Narayanan Kutty, C. Narayanan Kutty, and T. Pradeepkumar (2017). Fertigation - the key component of precision farming. *Journal of Tropical Agriculture*, 54(2):103-114.
- Tey, Y. S. and M. Brindal (2012). Factors influencing the adoption of precision agricultural technologies: a review for policy implications. *Precision Agriculture*, 13, 713-730.
- Tiamiyu, S., J. Akintola and M. Rahji (2009). Technology Adoption and Productivity Difference among Growers of New Rice for Africa in Savanna Zone of Nigeria.
- Tarjuelo, J. M., J. Montero, P. A. Carrion, F. T. Honrubia and M. A. Calvo (1999). Irrigation uniformity with medium size sprinklers part II: Influence of wind and other factors on water distribution. *Transactions of the ASAE*, 42, 677-689.

UCCE and CAS (University of California Cooperative Extension, San Diego County, and The California Avocado Society) (2012). *Avocado Production in California--A Cultural Handbook for Growers, Second Edition*. Book Two – Cultural Care.

USDA-Economic Research Service (2012). Beginning Farmers and Age Distribution of Farmers. Accessed on July 2, 2019 at <https://www.ers.usda.gov/topics/farm-economy/beginning-disadvantaged-farmers/beginning-farmers-and-age-distribution-of-farmers/>.

Wang, J., R. Mendelsohn, A. Dinar, and J. Huang (2010). How Chinese Farmers Change Crop Choice to Adapt to Climate Change. *Climate Change Economics*, 1(3) 167–185.

Yaron, D., A. Dinar and H. Voet (1992). Innovation on Family Farms: The Case of the Nazareth Region in Israel. *American J. of Agricultural Economics*, 74(2):361-370.

Table 1. Summary of Explanatory Variables

Determinant of Adoption	Predictor variables used in this study	Source	Hypotheses regarding impact on adoption (holding everything else constant)
Socio-economic characteristics	Operator age	Genius et al., 2014	Older growers full time growers with more years of experience, higher formal education, higher share of income from agriculture or of a given crop that is analyzed, complex organization types, or original ownership and larger operations and with higher cost of water will be adopters of technologies.
	Years of Experience	Tiamiyul, 2009	
	Formal Education	Robertson et al., 2012	
	Ownership Type	Campbell and Dinar, 1992	
	Cost of water	Feder and Umali., 1993	
	Share of Income from Farm (or Avocado)	Yaron et al., 1992	
Farm characteristics	Location of farm	Bryant et al., 2002	Growers in farms managing greater irrigation complexity will adopt technologies and management practices to help them cope with the difficulties of a more complex irrigation set up.
	Soil type, soil quality, topography	Bryant et al., 2002	
	Land tenure, full time operators	Fleischer et al., 2011	
	Farm size	Bohlen, Beal and Rogers, (1957)	
Informational	Use of cooperative extension	Genius et al., 2014; Tiamiyul, 2009; Bohlen, Beal and Rogers, (1957)	Growers who use cooperative extension and place a high level of importance on cooperative extension will adopt water conservation technologies.

Table 2. Distribution of Avocado Growers in the Sample

California County	Sample			California*	
	No. of Growers	Total Avocado (Acres)	Total Farm Land (Acres)	No. of Growers	Total Avocado (Acres)
Orange	3 (3)	94 (2)	1002	Unknown	Unknown
Riverside	15 (12)	734 (19)	836	Unknown	6127 (11)
San Diego	51 (41)	1036 (26)	1905	2000 (38)	20643 (38)
Santa Barbara	12 (10)	304 (8)	164	1000 (20)	5707 (11)
San Luis Obispo	7 (6)	151 (4)	2536	135 (4)	4214 (8)
Ventura	35 (28)	1580 (41)	4997	2000 (38)	17089 (32)
Total	123 (100)	3899 (100)	11440	5135 (100)	53780 (100)

*Approximately. California land and number of growers by county are from 2013 California Avocado Commission Report (2013) and personal interviews with UC Cooperative Extension agents.

In parenthesis are shares of county growers and total avocado acres of total California values.

Table 3. Distribution of Bundles in the Sample

Bundle	Number of Growers	Percent in Dataset
0	20	16.26
1	15	12.19
2	43	34.95
3	45	36.58
Total	123	$\cong 100$

Table 4. Descriptive Statistics of Variables Included in the Analysis

Variable	Obs.	Mean	Std. Dev.	Min	Max	
Water Audit	123	0.495	0.502	0	1	Technologies Adopted
Soil Moisture by Feel	123	0.528	0.501	0	1	
CIMIS	123	0.284	0.453	0	1	
Soil Moisture with Gypsum Block	123	0.024	0.154	0	1	
Soil Moisture with Tensiometer	123	0.317	0.467	0	1	
Calendar Irrigation	123	0.130	0.337	0	1	
Management of Tree Canopy	123	0.479	0.501	0	1	
Pressure Compensating Sprinkler	123	0.439	0.498	0	1	
Farm Acres	123	93.007	214.939	1	1100	Farm Characteristics
Avocado Acres	123	31.699	72.514	1	550	
Riverside County	123	0.122	0.329	0	1	
San Diego County	123	0.415	0.495	0	1	
Irrigation Complexity	123	0.405	1.013	0	8	
Agricultural water rate (\$/hcf) ^a	123	2.190	0.298	0	3.93	
Age of owner	123	61.910	11.502	27	88	Farmer Characteristics
Formal education	123	4.093	1.192	1	6	
Years' experience	123	19.830	14.223	1	58	
Share income from avocado	123	0.162	0.278	0	1	
Original Owner	123	0.422	0.496	0	1	
Use of Cooperative Extension	123	4.098	1.197	1	5	Informational Factors
Leaf Sampling	123	0.796	0.404	0	1	
Follow lab recommendation	123	1.861	1.147	0	3	
Test Irrigation Water	123	0.528	0.501	0	1	

^aNote: hcf = hundred of cubic feet. A term typically found in a grower's water bill. 1 cf = 0.0283 cubic meters.

Table 5. Logit regression models

Dependent Variable=Adoption as a binary choice of any saving technology or management practice.		
Variables	Logit Model 1	Logit Model 2
Owner Operated	0.259 (1.165)	-
Grow crops other than avocado?	-	-1.188* (0.739)
Farm Acres	0.00381 (0.00440)	-
Avocado acres	-	0.00248 (0.00635)
Riverside County	2.128** (1.510)	-
San Diego County	1.321 (0.851)	-
Cost of irrigation water	-	0.358 (0.339)
Irrigation Complexity	-7.08** (0.332)	-
Owner Age	-0.0814** (0.0323)	-0.068** (0.0297)
Education	-0.222 (0.297)	0.0163 (0.244)
Share income from avocado	15.44* (8.899)	9.180** (4.595)
Cooperative Extension Use	1.185*** (0.392)	1.454*** (0.464)
Constant	3.867 (2.513)	2.794 (2.279)
LR chi2(9)	58.55	50.99***
Prob>chi ²	0.000***	0.000***
Pseudo R ²	0.536	0.467
Log likelihood	-25.321	-29.109
Observations	123	123


Note: Standard errors in parentheses. *** p<0.01, ** p<0.05, * p<0.1

Table 6. Multinomial Logit Regression

Dependent Variable: Adoption of water saving technologies and management practices bundles as multinomial logit framework			
Variables	Bundle 1	Bundle 2	Bundle 3
Owner operated	0.332 (1.340)	0.0745 (1.380)	0.249 (1.229)
Farm acres	0.00368 (0.00530)	5.77 e-05 (0.00671)	0.00548 (0.00528)
Riverside County	2.794 (2.427)	4.989** (2.462)	5.302** (2.322)
San Diego County	0.0719 (1.042)	0.541 (1.202)	2.598*** (0.979)
Irrigation complexity	-0.277 (0.345)	-0.802 (0.506)	-1.163** (0.518)
Owner age	-0.0573 (0.0383)	-0.113** (0.0441)	-0.106*** (0.0363)
Education	-0.146 (0.333)	0.804** (0.389)	0.131 (0.344)
Share income from avocado	14.86* (8.938)	15.05* (9.005)	14.91* (8.914)
Cooperative extension use	1.016** (0.418)	0.867* (0.469)	1.415*** (0.421)
Constant	1.237 (3.031)	7.176** (3.338)	2.318 (2.773)
LR Chi ² (27)	102.230		
Prob>chi ²	0.000***		
Pseudo R ²	0.378		
Log likelihood	-83.965		
Number of observations (Bundle 0 with 20 Obs.)	15	43	45

Note: Standard errors in parentheses. *** p<0.01, ** p<0.05, * p<0.1

Figure 2: Multiple Correspondence Analysis (MCA) Coordinate Plot



AUXILIARY MATERIAL

Appendix 1: **Water Saving Technologies and Management Practices Used by California**

Avocado Growers

Avocado growers use a variety of irrigation technologies and management practices to sustain their yields. In the following we provide a description of the various technologies and management practices we encountered in our survey of growers in California.

Soil Moisture Monitoring

Careful water management is critical in the productivity of an avocado crop as overwatering or under-watering can significantly reduce yields (Ferreira et al., 2008; Gil et al., 2012; Kiggundu et al., 2012; Kozlowski, 1997). Determining when and how much to irrigate is difficult for a grower by visual monitoring or by hand sample. In our study, nearly 53% of growers in the sample use a method we identify as “By Feel”, a visual technique where growers take a sample of soil, if the soil doesn’t appear moist by sight or feel then they irrigate. It is impossible to determine the soil-water energy when a grower uses “By feel” techniques since soil water potential can only be measured with technology (Saxton and Rawls, 2006; Stagakis et al., 2012). Another popular method used by growers is to irrigate by calendar date (13% of growers in this study), which does not consider weather, soil conditions or water demand of the tree. Although, it’s important to note that irrigation by calendar date may be more likely in water districts where there are restricted allocations of water delivery and growers only have access to irrigation water on specific days.

Meteorological data to manage water applications

Weather stations are used in determining a crop’s water needs based on precipitation, temperature, wind and radiation combined with a crop coefficient to estimate evapotranspiration. In California, growers have access to a free web based evapotranspiration estimator, the California Irrigation

Management Information System (CIMIS) a system that was designed for irrigators to use their water resources more efficiently. CIMIS has 200 weather stations throughout California, growers can access weather data from a station nearby and, when combined with a crop coefficient, can be used to guide a specific crop needs. Twenty eight percent of the avocado growers in our sample use CIMIS as an irrigation calculator to determine when and how much water to apply. In general, irrigation calculators use daily values for evapotranspiration and rainfall to recommend weekly water inputs (Green et al. 2012). Though there are other types of irrigation software growers can use, in this study we only look at CIMIS as it is free to all California growers.

Canopy reduction

When faced with water shortages growers can choose from several management decisions to mitigate permanent damage to the trees. One type of extreme management method growers can utilize is the decision to stump trees, aggressively pruning the tree down to 4 to 7 feet from the ground. Stumping is used in order to save orchard trees during periods of drought because trees can be rejuvenated when water supply levels are revamped. When trees are stumped, the water requirement is drastically lowered due to a significant reduction of leaf area resulting in a lower loss of water from transpiration (Hofshi 2010; Schaffer et al., 2013). The drawback to this method is that the tree will not produce fruit for up to 5 years after rejuvenation, resulting in an economic hardship to the grower. Nearly 48% of the growers in our sample stump trees in their groves. Stumping, or canopy reduction, can also be used as a way to rejuvenate an aging orchard, however in this survey we asked if they stumped due to water shortage only.

Water Audits

Another water saving management decision includes improving irrigation performance by increasing an existing irrigation system's distribution uniformity (DU). DU, expressed as a

percentage, is an indication of how evenly distributed is the water application delivered to the crop. It accounts for differences in pressure, topography, and discharge coefficients from sprinklers or nozzles where optimal values for DU are considered 80% and above (Tarjuelo et al., 1999). The high uniformity of an irrigation system guarantees appropriate water delivery to the crop by reducing overwatering or under watering (Ascough and Kiker, 2002). Pereira, et al., (2002) found that improving irrigation efficiency, by increasing DU, is an important tool in water resource management because it reduces non-consumptive use of water and undesired water runoff. In most areas of California water audits (professional auditors) are offered for free by either the local water district or the resource conservation district. Nearly 50% of the growers in our sample used water audits.

Pressure Compensating Sprinkler

The aim of micro or drip irrigation systems are to minimize runoff and maximize water use to deliver water directly to the plant. However, it is important to consider distribution uniformity when designing an irrigation system to know how much water is being delivered to each tree. This is especially critical in orchards that have slopes or large areas of irrigation systems. Pressure compensating emitters are able to deliver a precise aliquot of water regardless of changes in pressure due to changes in terrain, such as slopes. Pressure compensating sprinklers are important in irrigation management because when used a grower know exactly how much water is being delivered to each plant. This results in a more uniform distribution where one tree isn't being overwatered and another underwatered.

Tensiometer


The most basic instruments for measuring plant available water are tensiometers and gypsum blocks (next section), both of which measure the “soil water potential”. The tensiometers are

installed to place the ceramic cup portion at a depth that matches with the root zone. PAW is the water fraction that over the range from field capacity (typically 12-18 centi-bars (cb)) up to 100 cb, where all plant available water has been depleted.

Gypsum Blocks

Another commonly used method for determining soil water potential is the gypsum blocks. These devices determine soil moisture by measuring the electrical resistance to current flow between electrodes that are embedded within a block of gypsum, or a similar material. The gypsum block allows moisture to move in and out as the soil becomes more saturated or dries out. When more moisture is absorbed by the block it lowers the resistance reading indicating a more saturated soil. The blocks are inexpensive and are easy to replace but require a data logger in order to get the readings. In addition, the blocks eventually dissolve and need to be replaced. As with tensiometers, gypsum blocks are somewhat slow to respond to rapid changes in soil moisture. They are the most useful for measuring the slow dry-down of soil over time, and thus are used to guide when irrigation should begin. On the other hand, the slow response time limits their utility for determining when to turn off the water, which can lead to overwatering when irrigation valves are directly controlled.

Appendix 2: KSOM of all adoption variables considered for use in the adoption bundles



Appendix 2 Legend	
Tile No.	Description
1	Clusters
2	Unified Distance Matrix
3	Test irrigation water
4	Test soil water
5	Measure total salts
6	Water audit? (Y/N)
7	District or groundwater for irrigation
8	Percent orchard on drip
9	Age of drip irrigation system
10	Percent orchard on micro irrigation
11	Age of micro irrigation

12	Use CIMIS? (Y/N)
13	By Feel
14	Gravimetric
15	Gypsum block
16	Tensiometer
17	Soil auger
18	Irrigate by calendar
19	Capacitance sensor
20	Dielectric sensor
21	Neutron probe
22	Pressure compensating sprinklers
23	Prune
24	Stump
25	Remove trees
26	Turn off water
27	Any tree management (23-26)

Appendix 3: The Survey Instrument - Adoption of water technologies and management practices by California avocado growers

FREE BOOK FOR YOUR PARTICIPATION IN THIS SURVEY! Recently published book on water policy in California "Managing California's Water, From Conflict to Reconciliation", 2011 (valued at \$35).

1. Where would you like your free book sent? Address

2. Please enter your email address if you want an electronic copy of the survey results

3. Total farm acreage under your ownership

4. How many acres of avocado

5. Total acres rented or leased for the 2012 year

6. Is avocado the only crop grown at your orchard?

Yes No

If no, what are the other types of crops are grown (please list top three)?

First crop: _____

Second crop: _____

Third crop: _____

Total acres of crops **other than avocado** grown at your orchard

What county is the orchard located in?

Los Angeles County Orange Riverside San Diego San Luis Obispo

Santa Barbara Ventura Other

Ownership type (select from list or fill in type):

Sole proprietorship: owned by one person

Partnering: two or more people share ownership

Corporations: shareholders elect a board of directors to oversee the major policies and decisions

LLC: type of hybrid business structure where owners are members

Other _____

Primary owners/managers age _____

Primary owners/managers Gender

Male Female

What was the highest level of education for the primary owner/manager?

	High school	Some college	A.A./A.S.	B.A./B.S.	Ma/PhD	Other
Education						

Years of experience growing avocado? _____

Years of experience growing crops other than avocado. _____

Are you the original owner?

Yes No

What year did you take ownership of the orchard? _____

Irrigation block acreage, shape, slope and soil texture. The factors that determine installation of irrigation technology.

Irrigation block acreage	Block shape Square <input type="checkbox"/>	Slope None = 0% Low = 0-5%	Soil texture: Sand Sandy loam
-----------------------------	--	----------------------------------	-------------------------------------

	Rectangular  Triangle  Irregular 	Med = 5-10 Mod = 10-15 Severe = <15%	Loam Clay loam Clay

Avocado tree characteristics such as variety, root stock, age, height, tree spacing, planting type.

# of trees	Variety and Rootstock	Avg. age of trees (yrs)	Avg. Height of trees (ft)	Tree spacing (ft) Ex. 20 X 20	1 = mound(berm) or 2 = native terrain

Orchard Management Characteristics

How often does a PCA visit the orchard?

Monthly Bi-annual Annually Not Applicable/Not using

How often do you treat for weed management (chemical or manual)?

Weekly Monthly Bi-annually Annually None

Fertilization strategy

Nitrogen Fertilization – Fertigation

Frequency (circle months)

Dec Jan Feb Mar Apr May Jun Jul Aug Sep Oct Nov

Hand application: What type(s) _____

Frequency (circle months)

Dec Jan Feb Mar Apr May Jun Jul Aug Sep Oct Nov

Aerial application: Material (s) _____

Frequency (circle months)

Dec Jan Feb Mar Apr May Jun Jul Aug Sep Oct Nov

Organic materials: Material (s) _____

Frequency (circle months)

Dec Jan Feb Mar Apr May Jun Jul Aug Sep Oct Nov

Do you take leaf samples?

Yes No

How often do you send leaf samples to a lab to get tested for nutrients?

Monthly Bi-annually Annually Never

If you send samples to a lab, how often do you follow their fertilization recommendations?

Always Sometimes Never Not applicable

How often do you test your irrigation water for total salts?

Monthly Bi-annually Annually Never

How often do you test your soil water for irrigation purposes?

Monthly Bi-annually Annually Never

Do you use CIMIS for soil water management?

Yes No

How often do you measure your soil for total salt (EC)?

Monthly Bi-annually Annually Never

What is your average soil EC in dS/m? (If known)

Environmental characteristics and Water delivery

Is there a privately owned weather station located at your orchard?

Yes No

How far away from your orchard is the nearest weather station that you are able to access?

>1 1-5 5-10 10-15 15-20 <20

What year did you last get a water audit on your orchard?

What is the water district of the orchard? Please fill in the boxes below.

	Water District 1	Water District 2
Water District Name		
Acres of your orchard serviced (ac)		
Annual water use for 2012 yr (ac/ft yr)		
Avg. Cost of water per month (\$/ac ft)		

Is water delivery service from your district available all year?

Yes No

Water delivery availability (select which one best describes your access to water):

- On demand - can be accessed at all times
- Rotation/schedule - limited water access to specific times or days
- Other

What is the lead time for water in hours?

Duration of Flows in hours?

What is the size of your water meter in inches? (If known)

Is there a restrictor on your water meter?

- Yes No

Type of water used for irrigation:

	District water	Ground water (well)	Recycled water	Trucked in water	Other
--	----------------	---------------------	----------------	------------------	-------

% used for irrigation					
If known fill in:					
ECw (dS/m)					
TDS (ppm)					
Chloride (ppm)					

Is there a water treatment system onsite?

Yes No

If there is a water treatment system onsite, what type do you use?

Reverse osmosis Ion exchange Distillation Other _____

Irrigation technologies and conservation practices

Fill in the percentage of each type of irrigation system installed and the average age of the system.

	Surface drip	Micro sprinklers	Other
% of grove with this type of system			
Age of system (yrs)			

How do you determine your orchard's soil-water status? Check all that apply.

CIMIS by feel(probe) gravimetric gypsum block tensiometer

auger method calendar capacitance probe Dielectric permittivity probe

neutron probe none other

Do you use pressure compensating emitters as part of your irrigation system?

Yes No

What is your typical irrigation schedule per season?

	Fall	Winter	Spring	Summer
Daily				
Weekly				
Bi-weekly				
Monthly				
Bi-monthly				
Other				

What is the application rate of your sprinklers in gallons per hour?

Is there a full time or dedicated irrigator employed at your orchard?

Yes No

At what frequency do you leach?

Daily Weekly Monthly Other _____ I don't leach

What is your % leaching fraction? (If known) _____

Orchard management structure

Is the orchard owner-operated? (If yes, skip next 3 questions)

Yes No

If No, who operates the orchards daily activities?

Management Company Farm Advisor Other

How often do you communicate with them?

Daily Weekly Monthly Bi-annually Annually

What method of communication do you use?

Phone Email In person Post mail Other

Rate the level of involvement from the management company or farm advisor?

Intense-makes all decisions

Moderate-makes most of the decisions

Medium-makes half the decisions

Low-limited decision making

Very low-makes very few decisions

Was there off farm employment by primary owner/manager in 2012?

Yes No

What was primary owners/managers % of income from avocado production in 2012?

Select the organization type that best fits your orchard management by circling the diagram that represents your orchard. Adapted from: Campbell and Dinar, 1993

1.Unified: Operated by a single individual who performs all	2.Cooperative Market Organization: The simplest level of horizontal task specialization
--	--

<p>the tasks</p> <div style="text-align: center; border: 1px solid black; width: fit-content; margin: 0 auto; padding: 5px;">Farm Production Unit</div>	<p>- the separation between workers and management</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Mechanics</div> <div style="font-size: 20px;">→</div> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Irrigators</div> <div style="font-size: 20px;">→</div> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Machine Operators</div> </div>
<p>3.Primary Hierarchy: Workers are distinguished from management as well as from each other. Two units: management and labor</p> <div style="text-align: center; margin-top: 20px;"> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 0 auto;">Management</div> <div style="font-size: 24px; margin: 5px auto;">↓</div> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 0 auto;">Production Unit</div> </div>	<p>4. Functional Hierarchy: Similar to simple but managers are also organized according to tasks. Irrigation managers perform distinct tasks from those performed by the machine shop manager.</p> <div style="text-align: center; margin-top: 20px;"> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 0 auto;">Management</div> <div style="font-size: 24px; margin: 5px auto;">↓</div> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Equipment Operators</div> <div style="font-size: 20px;">→</div> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Irrigators</div> <div style="font-size: 20px;">→</div> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Mechanics</div> </div> </div>
<p>5.Cooperative Market hierarchy: Similar to Type 2, Workers and managers form specialized units producing unique crops for particular markets</p> <div style="display: flex; justify-content: space-around; align-items: center; margin-top: 20px;"> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Lemons</div> <div style="font-size: 20px;">→</div> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Cherimoyas</div> <div style="font-size: 20px;">→</div> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Avocado</div> </div>	<p>6.Market Hierarchy: Workers are organized by markets, clients and locations</p> <div style="text-align: center; margin-top: 20px;"> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 0 auto;">Management</div> <div style="font-size: 24px; margin: 5px auto;">↓</div> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Lemons</div> <div style="font-size: 20px;">→</div> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Cherimoyas</div> <div style="font-size: 20px;">→</div> <div style="border: 1px solid black; padding: 5px; margin: 0 10px;">Avocado</div> </div> </div>
<p>Other: Please describe and draw diagram</p>	

Management practices that address water scarcity in your orchard.

What was your past management in response to a water shortage or low quality water at your orchard?

Type	% of orchard treated	#of acres treated
Selective Pruning		
Stumping		
Removal of trees		
Turned water off		
Other		
None		

How many months in advance do you make decisions about water conservation treatments?

Information gathering and communication

Where and how often do you obtain information on avocado production?

	Cooperative extension	CAC	Online sources	Journals/ books	Supplier/ Agents/ labs	Growers	Other
--	-----------------------	-----	----------------	-----------------	------------------------	---------	-------

Weekly							
Monthly							
Bi-annually							
Yearly							
Other							

What are the topics that you ask about from these sources?

	Cooperative extension	CAC	Online sources	Journals/ books	Supplier/ agents/ labs	Growers	Other
Fertilizer							
Irrigation							
Pest/disease							
Harvest							
Pollination							
Water policy							
Other							

Rate the level of importance that you place on the information collected on issues related to avocado production.

	Rate level of importance 1-5 1=low 5=high
--	--

Cooperative extension	
CAC	
Online sources	
Journals/ books	
Supplier/ Agents/ labs	
Growers	
Other	

Harvest Procedures

When do harvest events occur? (Circle corresponding months for type of picking)

Size pick: Dec Jan Feb Mar Apr May Jun Jul Aug Sep Oct
Nov

Strip: Dec Jan Feb Mar Apr May Jun Jul Aug Sep Oct
Nov

What is the distance from your orchard to the packing house in miles?

Almost done! One more page...

Yield for 2012 production.

Yield data for year 2012 entered in lbs per size picked for the entire orchard

Fruit Size	Total yield (lbs) for entire orchard
24	
28	
32	
36	
40	
48	
60	
70	
84	
96	
#2	
culls	

What was your number of theft events during the year 2012?

Was is your estimated amount of loss due to theft for 2012?

Thank You!

Thank you for taking our survey. Your response is very important to us and our research

Sincerely,